



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON A CYLINDER THE INTERSECTION OF WHICH WITH A SPHERE WILL DEVELOP INTO AN ELLIPSE.*

By D. N. LEHMER, University of California.

1. In his *Mathematical Recreations* published about 1674 Van Etten describes a method of drawing an "oval" with a pair of compasses leaving the distance between the points of the instrument unaltered. His scheme was to stretch the drawing paper around a cylinder. The curve thus obtained is not an ellipse but a transcendental curve whose equation is

$$y^2 + 4b^2 \sin^2 \frac{x}{2b} = a^2,$$

where b is the radius of the cylinder and a the distance between the points of the compasses. The curve is clearly the developed intersection of a sphere of radius a with a cylinder of radius b , the center of the sphere being on the surface of the cylinder.

2. The question suggests itself to find the cylinder such that the curve as obtained above will develop into an ellipse. Take the origin at the center of the sphere and let the y -axis be an element of the cylinder. Let a plane parallel to the xz -plane cut out on the cylinder a curve of length s . Then the equation of the ellipse in question will be

$$\frac{y^2}{a^2} + \frac{s^2}{b^2} = 1.$$

To find the nature of the cylinder we have then the equations

$$x^2 + y^2 + z^2 = a^2 \dots\dots (1),$$

$$\frac{y^2}{a^2} + \frac{s^2}{b^2} = 1 \dots\dots (2),$$

$$\frac{ds}{dx} = \sqrt{1 + \frac{dz^2}{ax^2}} \dots\dots (3).$$

From these it is easy to derive the differential equation connecting x and z :

$$a^2(x^2 + z^2) \left(1 + \frac{dz^2}{ax^2} \right) = b^2 \left(z \frac{dz}{ax} + x \right)^2.$$

Changing to polar coördinates and reducing,

*Read before the San Francisco Section of the American Mathematical Society.

$$\frac{d\rho}{d\theta} = \pm \frac{a\rho}{\sqrt{(b^2 - a^2)}}, \text{ or } \kappa\rho = e^{\pm [a/\sqrt{(b^2 - a^2)}]\theta}.$$

The cylinder has therefore for its right section a logarithmic spiral or rather two arranged in opposite directions.

3. This curious result may be stated in the following way. If an elliptical hole be cut out of a sheet of paper and the sheet wrapped about a sphere the diameter of which equals the minor axis of the ellipse the sheet wraps up into a cylinder a cross section of which is a logarithmic spiral. A model illustrating this relation between the ellipse and the spiral is easily constructed.

BERKELEY, April, 1904.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

200. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

No matter what value x be given, the *numerical* value of the expression $(x+2)/(2x^2+3x+6)$ can never exceed $\frac{1}{3}$.

II. Solution by G. W. GREENWOOD, M.A., Professor of Mathematics and Astronomy in McKendree College, Lebanon, Ill.

We know that $y+1/y \leq 2$.

$$\therefore z/2 + 2/z \leq 2. \quad 2z + 8/z \leq 8. \quad 2z + 8/z - 5 \leq 3. \quad \frac{2z^2 - 5z + 8}{z} \leq 3.$$

Now put $z=x+2$ and we get

$$\frac{2x^2+3x+6}{x+2} \leq 3; \text{ i. e., } \frac{x+2}{2x^2+3x+6} \geq \frac{1}{3}.$$

202. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons. W. Va.

Express in the form of radicals the roots of the equation

$$x^9 + 9mx^7 + 27m^2x^5 + 30m^3x^3 + 9mx + 2r = 0.$$

I. Solution by A. H. HOLMES, Brunswick, Maine.

Writing $x^3 + 3mx = y$ the equation becomes

$$y^3 + 3m^3y + 2r = 0 \dots\dots\dots (1).$$